

## **Appendix C: References**

- Axtell, J (2002) Minimum viable populations e-mail message dated November 21, 2002.
- BLM Colorado State Office. (1991). BLM Colorado State Office Wilderness Study Report, Statewide Overview, Record of Decision (ROD).
- BLM. (1991). Record of Decision: Vegetation Treatment on BLM Lands in Thirteen Western States.
- BLM White River Field Office (1999). Environmental Assessment #CO-017-99-93-EA: Twin Buttes Allotment Management Plan.
- Coates-Markle, L. (2002). Minimum viable populations e-mail message November 26, 2002.
- Coates-Markle, L (2002). Minimum viable populations e-mail message November 25, 2002.
- Couvet, D. (2002). Deleterious effects of restricted gene flow in fragmented populations. *Conservation Biology*. April, 2002. pages 369-376.
- Cothran E. Gus. (2002). Genetic Analysis of the West Douglas CO Feral Horse Herd. Department of Veterinary Science, University of Kentucky. Lexington, KY.
- Cothran, E.Gus; Wild horse and burro program research documents e-mail message. May 2, 2003
- Krueper, D. J. Bart and T. Rich (2002) Response of Vegetation and Breeding Birds to the Removal of Cattle on the San Pedro River, Arizona (U.S.A.). BLM unpublished manuscript.
- Lande and Barrowclough (1987); Effective population size, genetic variation and their use in population management; Cambridge University Press
- NTIS (National Technical Information Service), U.S. Department of Commerce. (1980). Wild and Free Roaming Horses and Burros: Current Knowledge and Recommended Research.
- Saccheri, I.M. and Koussaari, M. (1998). Inbreeding and extinction in a butterfly metapopulation; *Nature Magazine* #392; pages 491 – 494
- Singer, F. and Zeigenfuss, L. (2000); Genetic effective population size in the Pryor Mountain Wild Horse Herd; Colorado State University
- Thornhill, N. W. (1993). The natural history of inbreeding and outbreeding. Chicago University Press